

NDC 10565-001-32

4

0

0

2

SOLUTION OF HYDROGEN PEROXIDE 3% USP 10 VOLUME

3%

SEALED FOR YOUR PROTECTION IF SEAL IS BROKEN OR MISSING DO NOT USE FIRST AID ANTISEPTIC
Made in USA

32 FL. OZ. (946 mL)

Mfg. by Hydrox LABORATORIES 825-B Tollgate Road Elgin, IL 60123

Drug Facts

Active ingredient

Purpose

Hydrogen Peroxide (stabilized) 3%......First aid antiseptic/Oral debriding agent

USES • first aid to help prevent the risk of infection in minor cuts, scrapes and burns
• aids in the removal of philegin, mucous, or other secretions associated with occasional screenouth

Warnings For external use only

Do not use . in the eyes or apply over large areas of the body . longer than 1 week

Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns

Stop use and ask a doctor if • the condition persists or gets worse

· sore mouth symptoms do not improve in 7 days

imitation, pain, or redness persists or worsens
 swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

First aid antiseptic:

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- If bandaged, let dry first

Oral debriding agent (oral rinse): adults and children 2 years of age and over:

- · mix with an equal amount of water
- swish around in the mouth over the affected area for at least 1 minute and then spit out
- use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor
- . children under 12 years of age should be supervised in the use of this product
- · children under 2 years of age: consult a dentist or doctor

Other information Keep tightly closed in a cool dark place.
Do not shake bottle. Hold away from face when opening.

Inactive ingredient purified water

1-07-D0014

Rev.1